

## **APPLICATION OF PIMONIDAZOLE – NORMAL TISSUE**

### **2. Bacterial Granulomas, Fungal Infected Tissue**

1. Kempf VA, Lebedziejewski M, Alitalo K, et al. Activation of hypoxia-inducible factor-1 in bacillary angiomas: evidence for a role of hypoxia-inducible factor-1 in bacterial infections. *Circulation* 2005; 111: 1054-62.
2. Aly S, Wagner K, Keller C, et al. Oxygen status of lung granulomas in *Mycobacterium tuberculosis*-infected mice. *J Pathol* 2006; 210: 298-305.
3. Aly S, Laskay T, Mages J, Malzan A, Lang R, Ehlers S. Interferon-gamma-dependent mechanisms of mycobacteria-induced pulmonary immunopathology: the role of angiostasis and CXCR3-targeted chemokines for granuloma necrosis. *J Pathol* 2007; 212: 295-305.
4. Lenaerts AJ, Hoff D, Aly S, et al. Location of persisting mycobacteria in a Guinea pig model of tuberculosis revealed by r207910. *Antimicrob Agents Chemother* 2007; 51: 3338-45.
5. Degrossoli A, Bosetto MC, Lima CB, Giorgio S. Expression of hypoxia-inducible factor 1alpha in mononuclear phagocytes infected with *Leishmania amazonensis*. *Immunol Lett* 2007; 114: 119-25.
6. Echevarria M, Munoz-Cabello AM, Sanchez-Silva R, Toledo-Aral JJ, Lopez-Barneo J. Development of cytosolic hypoxia and hypoxia-inducible factor stabilization are facilitated by aquaporin-1 expression. *J Biol Chem* 2007; 282: 30207-15.
7. Klinkenberg LG, Sutherland LA, Bishai WR, Karakousis PC. Metronidazole Lacks Activity against *Mycobacterium tuberculosis* in an In Vivo Hypoxic Granuloma Model of Latency. *J Infect Dis* 2008; 198: 275-83.
8. Peyssonnaud C, Boutin AT, Zinkernagel AS, Datta V, Nizet V, Johnson RS. Critical role of HIF-1alpha in keratinocyte defense against bacterial infection. *J Invest Dermatol* 2008; 128: 1964-8.
9. Via LE, Lin PL, Ray SM, et al. Tuberculous granulomas are hypoxic in guinea pigs, rabbits, and nonhuman primates. *Infect Immun* 2008; 76: 2333-40.
10. Araujo AP, Frezza TF, Allegretti SM, Giorgio S. Hypoxia, hypoxia-inducible factor-1alpha and vascular endothelial growth factor in a murine model of *Schistosoma mansoni* infection. *Exp Mol Pathol* 2010; 89: 327-33.
11. Grahl N, Puttikamonkul S, Macdonald JM, et al. In vivo hypoxia and a fungal alcohol dehydrogenase influence the pathogenesis of invasive pulmonary aspergillosis. *PLoS Pathog* 2011; 7: e1002145.
12. Hempel C, Combes V, Hunt NH, Kurtzhals JA, Grau GE. CNS hypoxia is more pronounced in murine cerebral than noncerebral malaria and is reversed by erythropoietin. *Am J Pathol* 2011; 179: 1939-50.
13. Heng Y, Seah PG, Siew JY, et al. *Mycobacterium tuberculosis* infection induces hypoxic lung lesions in the rat. *Tuberculosis (Edinb)* 2011; 91: 339-41.

14. Kang YJ, Jo JO, Cho MK, Yu HS, Ock MS, Cha HJ. *Trichinella spiralis* infection induces angiogenic factor thymosin beta4 expression. *Vet Parasitol* 2011; 181: 222-8.
15. Thayil SM, Albin TA, Nazari H, et al. Local ischemia and increased expression of vascular endothelial growth factor following ocular dissemination of *Mycobacterium tuberculosis*. *PLoS One* 2011; 6: e28383.
16. Araujo AP, Arrais-Silva WW, Giorgio S. Infection by *Leishmania amazonensis* in mice: A potential model for chronic hypoxia. *Acta Histochem* 2012; 114: 797-804.
17. Driver ER, Ryan GJ, Hoff DR, et al. Evaluation of a mouse model of necrotic granuloma formation using C3HeB/FeJ mice for testing of drugs against *Mycobacterium tuberculosis*. *Antimicrob Agents Chemother* 2012; 56: 3181-95.
18. Harper J, Skerry C, Davis SL, et al. Mouse model of necrotic tuberculosis granulomas develops hypoxic lesions. *J Infect Dis* 2012; 205: 595-602.