

APPLICATION OF PIMONIDAZOLE – NORMAL TISSUE

4. Normal Tissues – Preclinical Studies

Bladder

1. Ghafar MA, Anastasiadis AG, Olsson LE, et al. Hypoxia and an angiogenic response in the partially obstructed rat bladder. *Lab Invest* 2002; 82: 903-9.
2. Ghafar MA, Shabsigh A, Chichester P, et al. Effects of chronic partial outlet obstruction on blood flow and oxygenation of the rat bladder. *J Urol* 2002; 167: 1508-12.
3. Levin RM, O'Connor LJ, Leggett RE, Whitbeck C, Chichester P. Focal hypoxia of the obstructed rabbit bladder wall correlates with intermediate decompensation. *Neurourol Urodyn* 2003; 22: 156-63.
4. Levin R, Chichester P, Levin S, Buttyan R. Role of angiogenesis in bladder response to partial outlet obstruction. *Scand J Urol Nephrol Suppl* 2004: 37-47.
5. Damaser MS, Whitbeck C, Chichester P, Levin RM. Effect of vaginal distension on blood flow and hypoxia of urogenital organs of the female rat. *J Appl Physiol* 2005; 98: 1884-90.
6. Lin AD, Mannikarottu A, Chaudhry A, et al. Protective effects of grape suspension on in vivo ischaemia/reperfusion of the rabbit bladder. *BJU Int* 2005; 96: 1397-402.
7. Badger WJ, Whitbeck C, Kogan B, Chichester P, Levin RM. The immediate effect of castration on female rabbit bladder blood flow and tissue oxygenation. *Urol Int* 2006; 76: 264-8.
8. Burgu B, Medina Ortiz WE, Pitera JE, Woolf AS, Wilcox DT. Vascular endothelial growth factor mediates hypoxic stimulated embryonic bladder growth in organ culture. *J Urol* 2007; 177: 1552-7.
9. Lin WY, Mannikarottu A, Li S, et al. Correlation of in vivo bladder blood flow measurements with tissue hypoxia. *World J Urol* 2009.
- b10. Morelli A, Filippi S, Comeglio P, et al. Acute vardenafil administration improves bladder oxygenation in spontaneously hypertensive rats. *J Sex Med* 2010; 7: 107-20.
11. Drzewiecki BA, Anumanthan G, Penn HA, et al. Modulation of the hypoxic response following partial bladder outlet obstruction. *J Urol* 2012; 188: 1549-54.

Bone

1. Dodd JS, Raleigh JA, Gross TS. Osteocyte hypoxia: a novel mechanotransduction pathway. *Am J Physiol Renal Physiol* 1999; 277: C598-602.
2. Davies Cd, Lundstrom LM, Frengen J, et al. Radiation improves the distribution and uptake of liposomal doxorubicin (caelyx) in human osteosarcoma xenografts. *Cancer Res* 2004; 64: 547-53.

3. Amarilio R, Viukov SV, Sharir A, Eshkar-Oren I, Johnson RS, Zelzer E. HIF1alpha regulation of Sox9 is necessary to maintain differentiation of hypoxic prechondrogenic cells during early skeletogenesis. *Development* 2007; 134: 3917-28.
4. Olmsted-Davis E, Gannon FH, Ozen M, et al. Hypoxic adipocytes pattern early heterotopic bone formation. *Am J Pathol* 2007; 170: 620-32.
5. Shafer J, Davis AR, Gannon FH, et al. Oxygen tension directs chondrogenic differentiation of myelo-monocytic progenitors during endochondral bone formation. *Tissue Eng* 2007; 13: 2011-9.
6. Genin O, Hasdai A, Shinder D, Pines M. Hypoxia, hypoxia-inducible factor-1alpha (HIF-1alpha), and heat-shock proteins in tibial dyschondroplasia. *Poult Sci* 2008; 87: 1556-64.
7. Kim HK, Bian H, Aya-ay J, Garces A, Morgan EF, Gilbert SR. Hypoxia and HIF-1alpha expression in the epiphyseal cartilage following ischemic injury to the immature femoral head. *Bone* 2009; 45: 280-8.
8. Fedorovich NE, Kuipers E, Gawlitta D, Dhert WJ, Alblas J. Scaffold porosity and oxygenation of printed hydrogel constructs affect functionality of embedded osteogenic progenitors. *Tissue Eng Part A* 2011; 17: 2473-86.
9. Criswell KA, Cook JC, Morse D, et al. Pregabalin induces hepatic hypoxia and increases endothelial cell proliferation in mice, a process inhibited by dietary vitamin E supplementation. *Toxicol Sci* 2012; 128: 42-56.
10. Donizetti-Oliveira C, Semedo P, Burgos-Silva M, et al. Adipose Tissue-Derived Stem Cell Treatment Prevents Renal Disease Progression. *Cell Transplant* 2012.

Bone Marrow

1. Asosingh K, De Raeve H, de Ridder M, et al. Role of the hypoxic bone marrow microenvironment in 5T2MM murine myeloma tumor progression. *Haematologica* 2005; 90: 810-7.
2. Levesque JP, Winkler IG, Hendy J, et al. Hematopoietic Progenitor Cell Mobilization Results in Hypoxia with Increased Hypoxia-Inducible Transcription Factor-1{alpha} and Vascular Endothelial Growth Factor A in Bone Marrow. *Stem Cells* 2007; 25: 1954-65.
3. Parmar K, Mauch P, Vergilio JA, Sackstein R, Down JD. Distribution of hematopoietic stem cells in the bone marrow according to regional hypoxia. *Proc Natl Acad Sci U S A* 2007; 104: 5431-6.
4. Brown AP, Chung EJ, Urlick ME, et al. Evaluation of the fullerene compound DF-1 as a radiation protector. *Radiat Oncol* 2010; 5: 34.
5. Hu J, Handisides DR, Van Valckenborgh E, et al. Targeting the multiple myeloma hypoxic niche with TH-302, a hypoxia-activated prodrug. *Blood* 2010; 116: 1524-7.
6. Zaleska K, Bruechner K, Baumann M, Zips D, Yaromina A. Tumour-infiltrating CD11b+ myelomonocytes and response to fractionated irradiation of human squamous cell carcinoma (hSCC) xenografts. *Radiother Oncol* 2011.

7. Jing D, Wobus M, Poitz DM, Bornhauser M, Ehninger G, Ordemann R. Oxygen tension plays a critical role in the hematopoietic microenvironment in vitro. *Haematologica* 2012; 97: 331-9.
8. Urao N, McKinney RD, Fukai T, Ushio-Fukai M. NADPH oxidase 2 regulates bone marrow microenvironment following hindlimb ischemia: role in reparative mobilization of progenitor cells. *Stem Cells* 2012; 30: 923-34.

Brain

1. Lee YM, Jeong CH, Koo SY, et al. Determination of hypoxic region by hypoxia marker in developing mouse embryos in vivo: a possible signal for vessel development. *Dev Dyn* 2001; 220: 175-86.
2. Forder JP, Munzenmaier DH, Greene AS. Angiogenic protection from focal ischemia with angiotensin II type 1 receptor blockade in the rat. *Am J Physiol Heart Circ Physiol* 2005; 288: H1989-96.
3. Morani A, Barros RP, Imamov O, et al. Lung dysfunction causes systemic hypoxia in estrogen receptor beta knockout (ERbeta^{-/-}) mice. *Proc Natl Acad Sci U S A* 2006; 103: 7165-9.
4. Nanka O, Valasek P, Dvorakova M, Grim M. Experimental hypoxia and embryonic angiogenesis. *Dev Dyn* 2006; 235: 723-33.
5. Noto T, Furuichi Y, Ishiye M, et al. Temporal and topographic profiles of tissue hypoxia following transient focal cerebral ischemia in rats. *J Vet Med Sci* 2006; 68: 803-7.
6. Kim YJ, Kang HH, Ahn JH, Chung JW. Hypoxic changes in the central nervous system of noise-exposed mice. *Acta Otolaryngol Suppl* 2007: 73-7.
7. Noto T, Furuichi Y, Ishiye M, et al. Tacrolimus (FK506) limits accumulation of granulocytes and platelets and protects against brain damage after transient focal cerebral ischemia in rat. *Biol Pharm Bull* 2007; 30: 313-7.
8. Ley EJ, Schemet J, Park R, et al. The in vivo effect of propranolol on cerebral perfusion and hypoxia after traumatic brain injury. *J Trauma* 2009; 66: 154-9; discussion 9-61.
9. Schelshorn DW, Schneider A, Kuschinsky W, et al. Expression of hemoglobin in rodent neurons. *J Cereb Blood Flow Metab* 2009; 29: 585-95.
10. Huang RQ, Cheng HL, Zhao XD, et al. Preliminary study on the effect of trauma-induced secondary cellular hypoxia in brain injury. *Neurosci Lett* 2010; 473: 22-7.
11. Leno-Duran E, Hatta K, Bianco J, et al. Fetal-placental hypoxia does not result from failure of spiral arterial modification in mice. *Placenta* 2010; 31: 731-7.
12. Hempel C, Combes V, Hunt NH, Kurtzhals JA, Grau GE. CNS hypoxia is more pronounced in murine cerebral than noncerebral malaria and is reversed by erythropoietin. *Am J Pathol* 2011; 179: 1939-50.
13. Hennings LJ, Flores R, Roberson PK, et al. Persistent penumbra in a rabbit stroke model: incidence and histologic characteristics. *Stroke Res Treat* 2011; 2011: 764830.

14. Del Bigio MR, Khan OH, da Silva Lopes L, Juliet PA. Cerebral white matter oxidation and nitrosylation in young rodents with kaolin-induced hydrocephalus. *J Neuropathol Exp Neurol* 2012; 71: 274-88.
15. Ito T, Funamoto K, Sato N, et al. Maternal undernutrition induces the expression of hypoxia-related genes in the fetal brain. *Tohoku J Exp Med* 2012; 226: 37-44.
16. Seiffge DJ, Lapina NE, Tsagogiorgas C, Theisinger B, Henning RH, Schilling L. Improvement of oxygen supply by an artificial carrier in combination with normobaric oxygenation decreases the volume of tissue hypoxia and tissue damage from transient focal cerebral ischemia. *Exp Neurol* 2012; 237: 18-25.
17. Yalcin O, Cabrales P. Increased hemoglobin O₂ affinity protects during acute hypoxia. *Am J Physiol Heart Circ Physiol* 2012; 303: H271-81.

Cartilage

1. Amarilio R, Viukov SV, Sharir A, Eshkar-Oren I, Johnson RS, Zelzer E. HIF1 α regulation of Sox9 is necessary to maintain differentiation of hypoxic prechondrogenic cells during early skeletogenesis. *Development* 2007; 134: 3917-28.
2. Shafer J, Davis AR, Gannon FH, et al. Oxygen tension directs chondrogenic differentiation of myelo-monocytic progenitors during endochondral bone formation. *Tissue Eng* 2007; 13: 2011-9.
3. Genin O, Hasdai A, Shinder D, Pines M. Hypoxia, hypoxia-inducible factor-1 α (HIF-1 α), and heat-shock proteins in tibial dyschondroplasia. *Poult Sci* 2008; 87: 1556-64.
4. Kim HK, Bian H, Aya-ay J, Garces A, Morgan EF, Gilbert SR. Hypoxia and HIF-1 α expression in the epiphyseal cartilage following ischemic injury to the immature femoral head. *Bone* 2009; 45: 280-8.

Digestive Tract

1. Cobb LM, Nolan J, Butler SA. Distribution of pimonidazole and RSU 1069 in tumour and normal tissues. *Br J Cancer* 1990; 62: 915-8.
2. Frangoulis M, Georgiou P, Chrisostomidis C, et al. Rat epigastric flap survival and VEGF expression after local copper application. *Plast Reconstr Surg* 2007; 119: 837-43.
3. Hernandez C, Santamatilde E, McCreath KJ, et al. Induction of trefoil factor (TFF)1, TFF2 and TFF3 by hypoxia is mediated by hypoxia inducible factor-1: implications for gastric mucosal healing. *Br J Pharmacol* 2009; 156: 262-72.
4. Hindryckx P, Waeytens A, Laukens D, et al. Absence of placental growth factor blocks dextran sodium sulfate-induced colonic mucosal angiogenesis, increases mucosal hypoxia and aggravates acute colonic injury. *Lab Invest* 2010; 90: 566-76.
5. Leno-Duran E, Hatta K, Bianco J, et al. Fetal-placental hypoxia does not result from failure of spiral arterial modification in mice. *Placenta* 2010; 31: 731-7.

6. Harris NR, Carter PR, Yadav AS, et al. Relationship between inflammation and tissue hypoxia in a mouse model of chronic colitis. *Inflamm Bowel Dis* 2011; 17: 742-6.
7. Hindryckx P, Devisscher L, Laukens D, Venken K, Peeters H, De Vos M. Intrarectal administration of oxygenated perfluorodecalin promotes healing of murine colitis by targeting inflammatory hypoxia. *Lab Invest* 2011; 91: 1266-76.
8. Yeh KY, Yeh M, Polk P, Glass J. Hypoxia-inducible factor-2alpha and iron absorptive gene expression in Belgrade rat intestine. *Am J Physiol Gastrointest Liver Physiol* 2011; 301: G82-90.

Ear

1. Donadieu E, Hamdi W, Deveze A, et al. Improved cryosections and specific immunohistochemical methods for detecting hypoxia in mouse and rat cochleae. *Acta Histochem* 2007; 109: 177-84.
2. Riva C, Donadieu E, Magnan J, Lavielle JP. Age-related hearing loss in CD/1 mice is associated to ROS formation and HIF target proteins up-regulation in the cochlea. *Exp Gerontol* 2007; 42: 327-36.
3. Cheeseman MT, Tyrer HE, Williams D, et al. HIF-VEGF pathways are critical for chronic otitis media in Junbo and Jeff mouse mutants. *PLoS Genet* 2011; 7: e1002336.

Embryo, Fetus

1. Lee YM, Jeong CH, Koo SY, et al. Determination of hypoxic region by hypoxia marker in developing mouse embryos in vivo: a possible signal for vessel development. *Dev Dyn* 2001; 220: 175-86.
2. Bassnett S, McNulty R. The effect of elevated intraocular oxygen on organelle degradation in the embryonic chicken lens. *J Exp Biol* 2003; 206: 4353-61.
3. Danielsson BR, Skold AC, Johansson A, Dillner B, Blomgren B. Teratogenicity by the hERG potassium channel blocking drug almokalant: use of hypoxia marker gives evidence for a hypoxia-related mechanism mediated via embryonic arrhythmia. *Toxicol Appl Pharmacol* 2003; 193: 168-76.
4. Freeburg PB, Robert B, St John PL, Abrahamson DR. Podocyte expression of hypoxia-inducible factor (HIF)-1 and HIF-2 during glomerular development. *J Am Soc Nephrol* 2003; 14: 927-38.
5. Freeburg PB, Abrahamson DR. Divergent expression patterns for hypoxia-inducible factor-1beta and aryl hydrocarbon receptor nuclear transporter-2 in developing kidney. *J Am Soc Nephrol* 2004; 15: 2569-78.
6. Danielsson BR, Johansson A, Danielsson C, Azarbayjani F, Blomgren B, Skold AC. Phenytoin teratogenicity: Hypoxia marker and effects on embryonic heart rhythm suggest an hERG-related mechanism. *Birth Defects Res A Clin Mol Teratol* 2005; 73: 146-53.
7. Nanka O, Valasek P, Dvorakova M, Grim M. Experimental hypoxia and embryonic angiogenesis. *Dev Dyn* 2006; 235: 723-33.

8. Webster WS, Howe AM, Abela D, Oakes DJ. The relationship between cleft lip, maxillary hypoplasia, hypoxia and phenytoin. *Curr Pharm Des* 2006; 12: 1431-48.
9. Amarilio R, Viukov SV, Sharir A, Eshkar-Oren I, Johnson RS, Zelzer E. HIF1alpha regulation of Sox9 is necessary to maintain differentiation of hypoxic prechondrogenic cells during early skeletogenesis. *Development* 2007; 134: 3917-28.
10. Burgu B, Medina Ortiz WE, Pitera JE, Woolf AS, Wilcox DT. Vascular endothelial growth factor mediates hypoxic stimulated embryonic bladder growth in organ culture. *J Urol* 2007; 177: 1552-7.
11. Oh C, Dong Y, Harman C, Mighty HE, Kopelman J, Thompson LP. Chronic hypoxia differentially increases glutathione content and gamma-glutamyl cysteine synthetase expression in fetal guinea pig organs. *Early Hum Dev* 2008; 84: 121-7.
12. Ream M, Ray AM, Chandra R, Chikaraishi DM. Early fetal hypoxia leads to growth restriction and myocardial thinning. *Am J Physiol Regul Integr Comp Physiol* 2008; 295: R583-95.
13. Nilsson MF, Danielsson C, Skold AC, et al. Improved methodology for identifying the teratogenic potential in early drug development of hERG channel blocking drugs. *Reprod Toxicol* 2010; 29: 156-63.
14. Shynlova O, Dorogin A, Lye SJ. Stretch-induced uterine myocyte differentiation during rat pregnancy: involvement of caspase activation. *Biol Reprod* 2010; 82: 1248-55.
15. Leno-Duran E, Hatta K, Bianco J, et al. Fetal-placental hypoxia does not result from failure of spiral arterial modification in mice. *Placenta* 2010; 31: 731-7.
16. Ito T, Funamoto K, Sato N, et al. Maternal undernutrition induces the expression of hypoxia-related genes in the fetal brain. *Tohoku J Exp Med* 2012; 226: 37-44.

Endometrium

1. Fan X, Krieg S, Kuo CJ, et al. VEGF blockade inhibits angiogenesis and reepithelialization of endometrium. *FASEB J* 2008.
2. Gillies RM, Robinson SP, McPhail LD, Carter ND, Murray JF. Immunohistochemical assessment of intrinsic and extrinsic markers of hypoxia in reproductive tissue: differential expression of HIF1alpha and HIF2alpha in rat oviduct and endometrium. *J Mol Histol* 2011; 42: 341-54.

Eye

1. Cobb LM, Nolan J, Butler SA. Distribution of pimonidazole and RSU 1069 in tumour and normal tissues. *Br J Cancer* 1990; 62: 915-8.
2. Gardiner TA, Gibson DS, de Gooyer TE, de la Cruz VF, McDonald DM, Stitt AW. Inhibition of Tumor Necrosis Factor- α Improves Physiological Angiogenesis and Reduces Pathological Neovascularization in Ischemic Retinopathy. *Am J Pathol* 2005; 166: 637-44.

3. Danylkova NO, Pomeranz HD, Alcala SR, McLoon LK. Histological and morphometric evaluation of transient retinal and optic nerve ischemia in rat. *Brain Res* 2006; 1096: 20-9.
4. de Gooyer TE, Stevenson KA, Humphries P, et al. Rod photoreceptor loss in Rho-/- mice reduces retinal hypoxia and hypoxia-regulated gene expression. *Invest Ophthalmol Vis Sci* 2006; 47: 5553-60.
5. de Gooyer TE, Stevenson KA, Humphries P, Simpson DA, Gardiner TA, Stitt AW. Retinopathy is reduced during experimental diabetes in a mouse model of outer retinal degeneration. *Invest Ophthalmol Vis Sci* 2006; 47: 5561-8.
6. Bartoli M, Al-Shabrawey M, Labazi M, et al. HMG-CoA reductase inhibitors (statins) prevents retinal neovascularization in a model of oxygen-induced retinopathy. *Invest Ophthalmol Vis Sci* 2008.
7. Saito Y, Uppal A, Byfield G, Budd S, Hartnett ME. Activated NAD(P)H Oxidase from Supplemental Oxygen Induces Neovascularization Independent of VEGF in Retinopathy of Prematurity Model. *Invest Ophthalmol Vis Sci* 2008; 49: 1591-8.
8. Holcombe DJ, Lengefeld N, Gole GA, Barnett NL. The effects of acute intraocular pressure elevation on rat retinal glutamate transport. *Acta Ophthalmol* 2008; 86: 408-14.
9. Bartoli M, Al-Shabrawey M, Labazi M, et al. HMG-CoA reductase inhibitors (statin) prevents retinal neovascularization in a model of oxygen-induced retinopathy. *Invest Ophthalmol Vis Sci* 2009; 50: 4934-40.
10. Chen J, Connor KM, Aderman CM, Willett KL, Aspegren OP, Smith LE. Suppression of retinal neovascularization by erythropoietin siRNA in a mouse model of proliferative retinopathy. *Invest Ophthalmol Vis Sci* 2009; 50: 1329-35.
11. Mowat FM, Luhmann UF, Smith AJ, et al. HIF-1alpha and HIF-2alpha are differentially activated in distinct cell populations in retinal ischaemia. *PLoS One* 2010; 5: e11103.
12. Wright WS, McElhatten RM, Messina JE, Harris NR. Hypoxia and the expression of HIF-1alpha and HIF-2alpha in the retina of streptozotocin-injected mice and rats. *Exp Eye Res* 2010; 90: 405-12.
13. Ly A, Yee P, Vessey KA, Phipps JA, Jobling AI, Fletcher EL. Early inner retinal astrocyte dysfunction during diabetes and development of hypoxia, retinal stress, and neuronal functional loss. *Invest Ophthalmol Vis Sci* 2011; 52: 9316-26.
14. Wright WS, McElhatten RM, Harris NR. Increase in retinal hypoxia-inducible factor-2alpha, but not hypoxia, early in the progression of diabetes in the rat. *Exp Eye Res* 2011.
15. Thayil SM, Albin TA, Nazari H, et al. Local ischemia and increased expression of vascular endothelial growth factor following ocular dissemination of *Mycobacterium tuberculosis*. *PLoS One* 2011; 6: e28383.
16. Wright WS, Yadav AS, McElhatten RM, Harris NR. Retinal blood flow abnormalities following six months of hyperglycemia in the Ins2(Akita) mouse. *Exp Eye Res* 2012; 98: 9-15.

Gonads

1. Vignozzi L, Filippi S, Morelli A, et al. Effect of chronic tadalafil administration on penile hypoxia induced by cavernous neurotomy in the rat. *J Sex Med* 2006; 3: 419-31.
2. Vignozzi L, Filippi S, Morelli A, et al. Cavernous neurotomy in the rat is associated with the onset of an overt condition of hypogonadism. *J Sex Med* 2009; 6: 1270-83.

Heart

1. Sinusas AJ. The potential for myocardial imaging with hypoxia markers. *Semin Nucl Med* 1999; 29: 330-8.
2. Lee YM, Jeong CH, Koo SY, et al. Determination of hypoxic region by hypoxia marker in developing mouse embryos in vivo: a possible signal for vessel development. *Dev Dyn* 2001; 220: 175-86.
3. Danielsson BR, Skold AC, Johansson A, Dillner B, Blomgren B. Teratogenicity by the hERG potassium channel blocking drug almokalant: use of hypoxia marker gives evidence for a hypoxia-related mechanism mediated via embryonic arrhythmia. *Toxicol Appl Pharmacol* 2003; 193: 168-76.
4. Danielsson BR, Johansson A, Danielsson C, Azarbayjani F, Blomgren B, Skold AC. Phenytoin teratogenicity: Hypoxia marker and effects on embryonic heart rhythm suggest an hERG-related mechanism. *Birth Defects Res A Clin Mol Teratol* 2005; 73: 146-53.
5. Forder JP, Munzenmaier DH, Greene AS. Angiogenic protection from focal ischemia with angiotensin II type 1 receptor blockade in the rat. *Am J Physiol Heart Circ Physiol* 2005; 288: H1989-96.
6. Cheema AN, Hong T, Nili N, et al. Adventitial microvessel formation after coronary stenting and the effects of SU11218, a tyrosine kinase inhibitor. *J Am Coll Cardiol* 2006; 47: 1067-75.
7. Morani A, Barros RP, Imamov O, et al. Lung dysfunction causes systemic hypoxia in estrogen receptor beta knockout (ERbeta^{-/-}) mice. *Proc Natl Acad Sci U S A* 2006; 103: 7165-9.
8. Zampino M, Yuzhakova M, Hansen J, et al. Sex-related dimorphic response of HIF-1 alpha expression in myocardial ischemia. *Am J Physiol Heart Circ Physiol* 2006; 291: H957-64.
9. Brown DA, MacLellan WR, Laks H, Dunn JC, Wu BM, Beygui RE. Analysis of oxygen transport in a diffusion-limited model of engineered heart tissue. *Biotechnol Bioeng* 2007; 97: 962-75.
10. Iheagwara KN, Thom SR, Deutschman CS, Levy RJ. Myocardial cytochrome oxidase activity is decreased following carbon monoxide exposure. *Biochim Biophys Acta* 2007.
11. Choi YH, Cowan DB, Nathan M, et al. Myocardial hypertrophy overrides the angiogenic response to hypoxia. *PLoS ONE* 2008; 3: e4042.

12. Lavine KJ, Kovacs A, Ornitz DM. Hedgehog signaling is critical for maintenance of the adult coronary vasculature in mice. *J Clin Invest* 2008; 118: 2404-14.
13. Ream M, Ray AM, Chandra R, Chikaraishi DM. Early fetal hypoxia leads to growth restriction and myocardial thinning. *Am J Physiol Regul Integr Comp Physiol* 2008; 295: R583-95.
14. Hennings LJ, Flores R, Roberson PK, et al. Persistent penumbra in a rabbit stroke model: incidence and histologic characteristics. *Stroke Res Treat* 2011; 2011: 764830.
15. Yalcin O, Cabrales P. Increased hemoglobin O₂ affinity protects during acute hypoxia. *Am J Physiol Heart Circ Physiol* 2012; 303: H271-81.

Kidney

1. Laurent F, Benard P, Canal P, Soula G. Autoradiographic distribution of [¹⁴C]-labelled pimonidazole in rhabdomyosarcoma-bearing rats and pigmented mice. *Cancer Chemother Pharmacol* 1988; 22: 308-15.
2. Cobb LM, Nolan J, Butler SA. Distribution of pimonidazole and RSU 1069 in tumour and normal tissues. *Br J Cancer* 1990; 62: 915-8.
3. Zhong Z, Arteel GE, Connor HD, et al. Cyclosporin A increases hypoxia and free radical production in rat kidneys: prevention by dietary glycine. *Am J Physiol* 1998; 275: F595-604.
4. Suga SI, Phillips MI, Ray PE, et al. Hypokalemia induces renal injury and alterations in vasoactive mediators that favor salt sensitivity. *Am J Physiol Renal Physiol* 2001; 281: F620-9.
5. Yin M, Zhong Z, Connor HD, et al. Protective effect of glycine on renal injury induced by ischemia- reperfusion in vivo. *Am J Physiol Renal Physiol* 2002; 282: F417-23.
6. Basile DP, Donohoe DL, Roethe K, Mattson DL. Chronic renal hypoxia after acute ischemic injury: effects of L-arginine on hypoxia and secondary damage. *Am J Physiol Renal Physiol* 2003; 284: F338-48.
7. Cachat F, Lange-Sperandio B, Chang AY, et al. Ureteral obstruction in neonatal mice elicits segment-specific tubular cell responses leading to nephron loss. *Kidney Int* 2003; 63: 564-75.
8. Freeburg PB, Robert B, St John PL, Abrahamson DR. Podocyte expression of hypoxia-inducible factor (HIF)-1 and HIF-2 during glomerular development. *J Am Soc Nephrol* 2003; 14: 927-38.
9. Freeburg PB, Abrahamson DR. Divergent expression patterns for hypoxia-inducible factor-1beta and aryl hydrocarbon receptor nuclear transporter-2 in developing kidney. *J Am Soc Nephrol* 2004; 15: 2569-78.
10. Manotham K, Tanaka T, Matsumoto M, et al. Evidence of tubular hypoxia in the early phase in the remnant kidney model. *J Am Soc Nephrol* 2004; 15: 1277-88.
11. Matsumoto M, Tanaka T, Yamamoto T, et al. Hypoperfusion of Peritubular Capillaries Induces Chronic Hypoxia before Progression of Tubulointerstitial Injury in a Progressive Model of Rat Glomerulonephritis. *J Am Soc Nephrol*, 2004; 15: 1574-81.

12. Samoszuk MK, Walter J, Mechetner E. Improved immunohistochemical method for detecting hypoxia gradients in mouse tissues and tumors. *J Histochem Cytochem* 2004; 52: 837-9.
13. Covington MD, Bayless KJ, Burghardt RC, Davis GE, Parrish AR. Ischemia-induced cleavage of cadherins in NRK cells: evidence for a role of metalloproteinases. *Am J Physiol Renal Physiol* 2005; 289: F280-8.
14. Rosenberger C, Heyman SN, Rosen S, et al. Up-regulation of HIF in experimental acute renal failure: Evidence for a protective transcriptional response to hypoxia. *Kidney Int* 2005; 67: 531-42.
15. Villanueva S, Cespedes C, Vio CP. Ischemic acute renal failure induces the expression of a wide range of nephrogenic proteins. *Am J Physiol Regul Integr Comp Physiol* 2005.
16. Wang PX, Sanders PW. Mechanism of hypertensive nephropathy in the Dahl/Rapp rat: a primary disorder of vascular smooth muscle. *Am J Physiol Renal Physiol* 2005; 288: F236-42.
17. Goldfarb M, Rosenberger C, Abassi Z, et al. Acute-on-chronic renal failure in the rat: functional compensation and hypoxia tolerance. *Am J Nephrol* 2006; 26: 22-33.
18. Morani A, Barros RP, Imamov O, et al. Lung dysfunction causes systemic hypoxia in estrogen receptor beta knockout (ERbeta^{-/-}) mice. *Proc Natl Acad Sci U S A* 2006; 103: 7165-9.
19. Rosenberger C, Rosen S, Shina A, et al. Hypoxia-inducible factors and tubular cell survival in isolated perfused kidneys. *Kidney Int* 2006; 70: 60-70.
20. Tanaka T, Kato H, Kojima I, et al. Hypoxia and expression of hypoxia-inducible factor in the aging kidney. *J Gerontol A Biol Sci Med Sci* 2006; 61: 795-805.
21. Villanueva S, Cespedes C, Vio CP. Ischemic acute renal failure induces the expression of a wide range of nephrogenic proteins. *Am J Physiol Regul Integr Comp Physiol* 2006; 290: R861-70.
22. Bernhardt WM, Wiesener MS, Weidemann A, et al. Involvement of hypoxia-inducible transcription factors in polycystic kidney disease. *Am J Pathol* 2007; 170: 830-42.
23. Higgins DF, Kimura K, Bernhardt WM, et al. Hypoxia promotes fibrogenesis in vivo via HIF-1 stimulation of epithelial-to-mesenchymal transition. *J Clin Invest* 2007; 117: 3810-20.
24. Imamura R, Moriyama T, Isaka Y, et al. Erythropoietin protects the kidneys against ischemia reperfusion injury by activating hypoxia inducible factor-1alpha. *Transplantation* 2007; 83: 1371-9.
25. Efrati S, Berman S, Ben Aharon G, Siman-Tov Y, Averbukh Z, Weissgarten J. Application of normobaric hyperoxia therapy for amelioration of haemorrhagic shock-induced acute renal failure. *Nephrol Dial Transplant* 2008; 23: 2213-22.
26. Heyman SN, Khamaisi M, Rosen S, Rosenberger C. Renal Parenchymal Hypoxia, Hypoxia Response and the Progression of Chronic Kidney Disease. *Am J Nephrol* 2008; 28: 998-1006.
27. Oh C, Dong Y, Harman C, Mighty HE, Kopelman J, Thompson LP. Chronic hypoxia differentially increases glutathione content and gamma-glutamyl cysteine synthetase expression in fetal guinea pig organs. *Early Hum Dev* 2008; 84: 121-7.

28. Rosenberger C, Goldfarb M, Shina A, et al. Evidence for sustained renal hypoxia and transient hypoxia adaptation in experimental rhabdomyolysis-induced acute kidney injury. *Nephrol Dial Transplant* 2008; 23: 1135-43.
29. Rosenberger C, Khamaisi M, Abassi Z, et al. Adaptation to hypoxia in the diabetic rat kidney. *Kidney Int* 2008; 73: 34-42.
30. Stoessel A, Paliege A, Theilig F, et al. Indolent course of tubulointerstitial disease in a mouse model of subpressor, low-dose nitric oxide synthase inhibition. *Am J Physiol Renal Physiol* 2008; 295: F717-25.
31. Wiggins KJ, Tiau V, Zhang Y, Gilbert RE, Langham RG, Kelly DJ. Perindopril attenuates tubular hypoxia and inflammation in an experimental model of diabetic nephropathy in transgenic Ren-2 rats. *Nephrology (Carlton)* 2008; 13: 721-9.
32. Fujimoto S, Satoh M, Nagasu H, Horike H, Sasaki T, Kashihara N. Azelnidipine exerts renoprotective effects by improvement of renal microcirculation in angiotensin II infusion rats. *Nephrol Dial Transplant* 2009.
33. Rosenberger C, Rosen S, Paliege A, Heyman SN. Pimonidazole adduct immunohistochemistry in the rat kidney: detection of tissue hypoxia. *Methods Mol Biol* 2009; 466: 161-74.
34. Shibata R, Ueda S, Yamagishi S, et al. Involvement of asymmetric dimethylarginine (ADMA) in tubulointerstitial ischaemia in the early phase of diabetic nephropathy. *Nephrol Dial Transplant* 2009; 24: 1162-9.
35. Palm F, Nangaku M, Fasching A, et al. Uremia induces abnormal oxygen consumption in tubules and aggravates chronic hypoxia of the kidney via oxidative stress. *Am J Physiol Renal Physiol* 2010; 299: F380-6.
36. Prasad P, Li LP, Halter S, Cabray J, Ye M, Battle D. Evaluation of renal hypoxia in diabetic mice by BOLD MRI. *Invest Radiol* 2010; 45: 819-22.
37. Yim HE, Kim JH, Yoo KH, et al. Spironolactone, but not enalapril, potentiates hypoxia-inducible factor-1 alpha and Ets-1 expression in newborn rat kidney. *J Physiol Pharmacol* 2010; 61: 73-81.
38. Takiyama Y, Harumi T, Watanabe J, et al. Tubular injury in a rat model of type 2 diabetes is prevented by metformin: a possible role of HIF-1alpha expression and oxygen metabolism. *Diabetes* 2011; 60: 981-92.
39. Yim HE, Kim JH, Yoo KH, Bae IS, Hong YS, Lee JW. Spironolactone and enalapril differentially up-regulate the expression of VEGF and heme oxygenase-1 in the neonatal rat kidney. *Pediatr Res* 2011; 69: 378-83.
40. Ding A, Kalaignanasundaram P, Ricardo SD, et al. Chronic treatment with tempol does not significantly ameliorate renal tissue hypoxia or disease progression in a rodent model of polycystic kidney disease. *Clin Exp Pharmacol Physiol* 2012.
41. Doi K, Katagiri D, Negishi K, et al. Mild elevation of urinary biomarkers in prerenal acute kidney injury. *Kidney Int* 2012.
42. Donizetti-Oliveira C, Semedo P, Burgos-Silva M, et al. Adipose Tissue-Derived Stem Cell Treatment Prevents Renal Disease Progression. *Cell Transplant* 2012.
43. Dupuy V, Mayeur N, Buleon M, et al. Type 2 diabetes mellitus in mice aggravates the renal impact of hemorrhagic shock. *Shock* 2012; 38: 351-5.
44. Nakazawa Y, Nishino T, Obata Y, et al. Recombinant human erythropoietin attenuates renal tubulointerstitial injury in murine adriamycin-induced nephropathy. *J Nephrol* 2012.

Liver

1. Laurent F, Benard P, Canal P, Soula G. Autoradiographic distribution of [14C]-labelled pimonidazole in rhabdomyosarcoma-bearing rats and pigmented mice. *Cancer Chemother Pharmacol* 1988; 22: 308-15.
2. Cobb LM, Nolan J, Butler SA. Distribution of pimonidazole and RSU 1069 in tumour and normal tissues. *Br J Cancer* 1990; 62: 915-8.
3. Arteel GE, Thurman RG, Yates JM, Raleigh JA. Evidence that hypoxia markers detect oxygen gradients in liver: pimonidazole and retrograde perfusion of rat liver. *Br J Cancer* 1995; 72: 889-95.
4. Arteel GE, Raleigh JA, Bradford BU, Thurman RG. Acute alcohol produces hypoxia directly in rat liver tissue in vivo: role of Kupffer cells. *Am J Physiol* 1996; 271: G494-500.
5. Arteel GE, Iimuro Y, Yin M, Raleigh JA, Thurman RG. Chronic enteral ethanol treatment causes hypoxia in rat liver tissue in vivo. *Hepatology* 1997; 25: 920-6.
6. Stachlewitz RF, Arteel GE, Raleigh JA, Connor HD, Mason RP, Thurman RG. Development and characterization of a new model of tacrine-induced hepatotoxicity: role of the sympathetic nervous system and hypoxia- reoxygenation. *J Pharmacol Exp Ther* 1997; 282: 1591-9.
7. Arteel GE, Thurman RG, Raleigh JA. Reductive metabolism of the hypoxia marker pimonidazole is regulated by oxygen tension independent of the pyridine nucleotide redox state. *Eur J Biochem* 1998; 253: 743-50.
8. Thurman RG. II. Alcoholic liver injury involves activation of Kupffer cells by endotoxin. *Am J Physiol* 1998; 275: G605-11.
9. Thurman RG, Bradford BU, Iimuro Y, et al. The role of gut-derived bacterial toxins and free radicals in alcohol- induced liver injury. *J Gastroenterol Hepatol* 1998; 13 Suppl: S39-50.
10. Rosmorduc O, Wendum D, Corpechot C, et al. Hepatocellular hypoxia-induced vascular endothelial growth factor expression and angiogenesis in experimental biliary cirrhosis. *Am J Pathol* 1999; 155: 1065-73.
11. Schemmer P, Bradford BU, Rose ML, et al. Intravenous glycine improves survival in rat liver transplantation. *Am J Physiol* 1999; 276: G924-32.
12. Schemmer P, Bunzendahl H, Raleigh JA, Thurman RG. Graft survival is improved by hepatic denervation before organ harvesting. *Transplantation* 1999; 67: 1301-7.
13. Schemmer P, Connor HD, Arteel GE, et al. Reperfusion injury in livers due to gentle in situ organ manipulation during harvest involves hypoxia and free radicals. *J Pharmacol Exp Ther* 1999; 290: 235-40.
14. Zhong Z, Arteel GE, Connor HD, et al. Binge drinking disturbs hepatic microcirculation after transplantation: prevention with free radical scavengers. *J Pharmacol Exp Ther* 1999; 290: 611-20.
15. Thurman RG. Sex-related liver injury due to alcohol involves activation of Kupffer cells by endotoxin. *Can J Gastroenterol* 2000; 14 Suppl D: 129D-35D.

16. Schemmer P, Enomoto N, Bradford BU, et al. Activated Kupffer cells cause a hypermetabolic state after gentle in situ manipulation of liver in rats. *Am J Physiol Gastrointest Liver Physiol* 2001; 280: G1076-82.
17. Schemmer P, Enomoto N, Bradford BU, Bunzendahl H, Raleigh JA, Thurman RG. Autonomic nervous system and gut-derived endotoxin: involvement in activation of kupffer cells after in situ organ manipulation. *World J Surg* 2001; 25: 399-406.
18. Zhong Z, Li X, Yamashina S, et al. Cyclosporin A causes a hypermetabolic state and hypoxia in the liver: prevention by dietary glycine. *J Pharmacol Exp Ther* 2001; 299: 858-65.
19. Bardag-Gorce F, French BA, Li J, et al. The importance of cycling of blood alcohol levels in the pathogenesis of experimental alcoholic liver disease in rats. *Gastroenterology* 2002; 123: 325-35.
20. Corpechot C, Barbu V, Wendum D, et al. Hepatocyte growth factor and c-Met inhibition by hepatic cell hypoxia: a potential mechanism for liver regeneration failure in experimental cirrhosis. *Am J Pathol* 2002; 160: 613-20.
21. Copple BL, Rondelli CM, Maddox JF, Hoglen NC, Ganey PE, Roth RA. Modes of cell death in rat liver after monocrotaline exposure. *Toxicol Sci* 2004; 77: 172-82.
22. Jeong WI, Do SH, Yun HS, et al. Hypoxia potentiates transforming growth factor-beta expression of hepatocyte during the cirrhotic condition in rat liver. *Liver Int* 2004; 24: 658-68.
23. Samoszuk MK, Walter J, Mechetner E. Improved immunohistochemical method for detecting hypoxia gradients in mouse tissues and tumors. *J Histochem Cytochem* 2004; 52: 837-9.
24. Forder JP, Munzenmaier DH, Greene AS. Angiogenic protection from focal ischemia with angiotensin II type 1 receptor blockade in the rat. *Am J Physiol Heart Circ Physiol* 2005; 288: H1989-96.
25. Luyendyk JP, Shaw PJ, Green CD, Maddox JF, Ganey PE, Roth RA. Coagulation-mediated hypoxia and neutrophil-dependent hepatic injury in rats given lipopolysaccharide and ranitidine. *J Pharmacol Exp Ther* 2005; 314: 1023-31.
26. Paizis G, Tikellis C, Cooper ME, et al. Chronic liver injury in rats and humans upregulates the novel enzyme angiotensin converting enzyme 2. *Gut* 2005; 54: 1790-6.
27. Copple BL, Roth RA, Ganey PE. Anticoagulation and inhibition of nitric oxide synthase influence hepatic hypoxia after monocrotaline exposure. *Toxicology* 2006; 225: 128-37.
28. Khan Z, Michalopoulos GK, Stolz DB. Peroxisomal localization of hypoxia-inducible factors and hypoxia-inducible factor regulatory hydroxylases in primary rat hepatocytes exposed to hypoxia-reoxygenation. *Am J Pathol* 2006; 169: 1251-69.
29. Morani A, Barros RP, Imamov O, et al. Lung dysfunction causes systemic hypoxia in estrogen receptor beta knockout (ERbeta^{-/-}) mice. *Proc Natl Acad Sci U S A* 2006; 103: 7165-9.
30. Tanaka H, Yamamoto M, Hashimoto N, et al. Hypoxia-independent overexpression of hypoxia-inducible factor 1alpha as an early change in mouse hepatocarcinogenesis. *Cancer Res* 2006; 66: 11263-70.

31. Cheluvappa R, Hilmer SN, Kwun SY, Cogger VC, DG LEC. Effects of old age on hepatocyte oxygenation. *Ann N Y Acad Sci* 2007; 1114: 88-92.
32. Cheluvappa R, Hilmer SN, Kwun SY, et al. The effect of old age on liver oxygenation and the hepatic expression of VEGF and VEGFR2. *Exp Gerontol* 2007; 42: 1012-9.
33. Deng X, Luyendyk JP, Zou W, et al. Neutrophil interaction with the hemostatic system contributes to liver injury in rats cotreated with lipopolysaccharide and ranitidine. *J Pharmacol Exp Ther* 2007; 322: 852-61.
34. Plock J, Frese S, Keogh A, et al. Activation of non-ischemic, hypoxia-inducible signalling pathways up-regulate cytoprotective genes in the murine liver. *J Hepatol* 2007; 47: 538-45.
35. Terada N, Ohno N, Saitoh S, Ohno S. Immunohistochemical detection of hypoxia in mouse liver tissues treated with pimonidazole using "in vivo cryotechnique". *Histochem Cell Biol* 2007; 128: 253-61.
36. Tugues S, Fernandez-Varo G, Munoz-Luque J, et al. Antiangiogenic treatment with sunitinib ameliorates inflammatory infiltrate, fibrosis, and portal pressure in cirrhotic rats. *Hepatology* 2007; 46: 1919-26.
37. Oh C, Dong Y, Harman C, Mighty HE, Kopelman J, Thompson LP. Chronic hypoxia differentially increases glutathione content and gamma-glutamyl cysteine synthetase expression in fetal guinea pig organs. *Early Hum Dev* 2008; 84: 121-7.
38. Riedl CC, Brader P, Zanzonico PB, et al. Imaging hypoxia in orthotopic rat liver tumors with iodine 124-labeled iodoazomycin galactopyranoside PET. *Radiology* 2008; 248: 561-70.
39. Zhao D, Najbauer J, Garcia E, et al. Neural stem cell tropism to glioma: critical role of tumor hypoxia. *Mol Cancer Res* 2008; 6: 1819-29.
40. Kao YH, Jawan B, Goto S, et al. Serum factors potentiate hypoxia-induced hepatotoxicity in vitro through increasing transforming growth factor-beta1 activation and release. *Cytokine* 2009; 47: 11-22.
41. Rosenberger C, Rosen S, Paliege A, Heyman SN. Pimonidazole adduct immunohistochemistry in the rat kidney: detection of tissue hypoxia. *Methods Mol Biol* 2009; 466: 161-74.
42. Schmeding M, Hunold G, Ariyakhagorn V, et al. Erythropoietin reduces ischemia-reperfusion injury after liver transplantation in rats. *Transpl Int* 2009; 22: 738-46.
43. Xia CY, Li L, Liu HM, Cong WM. High expression of angiotensin-converting enzyme and angiotensin-converting enzyme 2 in preservation injury after liver transplantation in rats. *Hepatol Res* 2009.
44. Araujo AP, Frezza TF, Allegretti SM, Giorgio S. Hypoxia, hypoxia-inducible factor-1alpha and vascular endothelial growth factor in a murine model of *Schistosoma mansoni* infection. *Exp Mol Pathol* 2010; 89: 327-33.
45. Schmeding M, Rademacher S, Boas-Knoop S, et al. rHuEpo reduces ischemia-reperfusion injury and improves survival after transplantation of fatty livers in rats. *Transplantation* 2010; 89: 161-8.
46. Chaudhuri S, McCullough SS, Hennings L, et al. Acetaminophen hepatotoxicity and HIF-1alpha induction in acetaminophen toxicity in mice occurs without hypoxia. *Toxicol Appl Pharmacol* 2011; 252: 211-20.

47. Criswell KA, Cook JC, Morse D, et al. Pregabalin induces hepatic hypoxia and increases endothelial cell proliferation in mice, a process inhibited by dietary vitamin E supplementation. *Toxicol Sci* 2012; 128: 42-56.

Lymph

1. Ohta A, Diwanji R, Kini R, Subramanian M, Sitkovsky M. In vivo T cell activation in lymphoid tissues is inhibited in the oxygen-poor microenvironment. *Front Immunol* 2011; 2: 27.
2. Zampell JC, Yan A, Avraham T, Daluvoy S, Weitman ES, Mehrara BJ. HIF-1alpha coordinates lymphangiogenesis during wound healing and in response to inflammation. *FASEB J* 2012; 26: 1027-39.

Ovary

1. Morani A, Barros RP, Imamov O, et al. Lung dysfunction causes systemic hypoxia in estrogen receptor beta knockout (ERbeta^{-/-}) mice. *Proc Natl Acad Sci U S A* 2006; 103: 7165-9.
2. Gillies RM, Robinson SP, McPhail LD, Carter ND, Murray JF. Immunohistochemical assessment of intrinsic and extrinsic markers of hypoxia in reproductive tissue: differential expression of HIF1alpha and HIF2alpha in rat oviduct and endometrium. *J Mol Histol* 2011; 42: 341-54.

Pancreas

1. Laurent F, Benard P, Canal P, Soula G. Autoradiographic distribution of [14C]-labelled pimonidazole in rhabdomyosarcoma-bearing rats and pigmented mice. *Cancer Chemother Pharmacol* 1988; 22: 308-15.
2. McKim SE, Uesugi T, Raleigh JA, McClain CJ, Arteel GE. Chronic intragastric alcohol exposure causes hypoxia and oxidative stress in the rat pancreas. *Arch Biochem Biophys* 2003; 417: 34-43.
3. Olsson R, Carlsson PO. A low-oxygenated subpopulation of pancreatic islets constitutes a functional reserve of endocrine cells. *Diabetes* 2011; 60: 2068-75.
4. Olsson R, Olerud J, Pettersson U, Carlsson PO. Increased numbers of low-oxygenated pancreatic islets after intraportal islet transplantation. *Diabetes* 2011; 60: 2350-3.
5. Sato Y, Endo H, Okuyama H, et al. Cellular hypoxia of pancreatic beta-cells due to high levels of oxygen consumption for insulin secretion in vitro. *J Biol Chem* 2011; 286: 12524-32.
6. Bensellam M, Duvillie B, Rybachuk G, et al. Glucose-induced O₂ consumption activates hypoxia inducible factors 1 and 2 in rat insulin-secreting pancreatic beta-cells. *PLoS One* 2012; 7: e29807.

Penis

1. Vignozzi L, Filippi S, Morelli A, et al. Effect of chronic tadalafil administration on penile hypoxia induced by cavernous neurotomy in the rat. *J Sex Med* 2006; 3: 419-31.

Peritoneum

1. Matsuzaki S, Jardon K, Maleysson E, et al. Carbon dioxide pneumoperitoneum, intraperitoneal pressure, and peritoneal tissue hypoxia: a mouse study with controlled respiratory support. *Surg Endosc* 2010; 24: 2871-80.

Placenta

1. Choy MY, Leung TN, Leung PS, Lau TK. First Trimester Human Trophoblast Production Of Placental Corticotrophin-Releasing Hormone (CRH) Is Unresponsive To Hypoxia In-Vitro. *Early Pregnancy: Biology and Medicine* 2003; VI: 235-47.
2. Pringle KG, Kind KL, Thompson JG, Roberts CT. Complex interactions between hypoxia inducible factors, insulin-like growth factor-II and oxygen in early murine trophoblasts. *Placenta* 2007; 28: 1147-57.
3. Leno-Duran E, Hatta K, Bianco J, et al. Fetal-placental hypoxia does not result from failure of spiral arterial modification in mice. *Placenta* 2010; 31: 731-7.
4. Kulandavelu S, Whiteley KJ, Qu D, Mu J, Bainbridge SA, Adamson SL. Endothelial nitric oxide synthase deficiency reduces uterine blood flow, spiral artery elongation, and placental oxygenation in pregnant mice. *Hypertension* 2012; 60: 231-8.

Prostate

1. Shabsigh A, Ghafar MA, de la Taille A, et al. Biomarker analysis demonstrates a hypoxic environment in the castrated rat ventral prostate gland. *J Cell Biochem* 2001; 81: 437-44.
2. Morani A, Barros RP, Imamov O, et al. Lung dysfunction causes systemic hypoxia in estrogen receptor beta knockout (ERbeta^{-/-}) mice. *Proc Natl Acad Sci U S A* 2006; 103: 7165-9.
3. Rudolfsson SH, Bergh A. Testosterone-stimulated growth of the rat prostate may be driven by tissue hypoxia and hypoxia-inducible factor-1alpha. *J Endocrinol* 2008; 196: 11-9.
4. Vignozzi L, Filippi S, Morelli A, et al. Cavernous neurotomy in the rat is associated with the onset of an overt condition of hypogonadism. *J Sex Med* 2009; 6: 1270-83.

Rectum

1. Greijer AE, Delis-van Diemen PM, Fijneman RJ, et al. Presence of HIF-1 and related genes in normal mucosa, adenomas and carcinomas of the colorectum. *Virchows Arch* 2008; 452: 535-44.

Skin

1. Cobb LM, Nolan J, Butler SA. Distribution of pimonidazole and RSU 1069 in tumour and normal tissues. *Br J Cancer* 1990; 62: 915-8.
2. Haroon ZA, Raleigh JA, Greenberg CS, Dewhirst MW. Early wound healing exhibits cytokine surge without evidence of hypoxia. *Ann Surg* 2000; 231: 137-47.
3. Albina JE, Mastrofrancesco B, Vessella JA, Louis CA, Henry WL, Jr., Reichner JS. HIF-1 expression in healing wounds: HIF-1alpha induction in primary inflammatory cells by TNF-alpha. *Am J Physiol Cell Physiol* 2001; 281: C1971-7.
4. Lokmic Z, Darby IA, Thompson EW, Mitchell GM. Time course analysis of hypoxia, granulation tissue and blood vessel growth, and remodeling in healing rat cutaneous incisional primary intention wounds. *Wound Repair Regen* 2006; 14: 277-88.
5. Erba P, Ogawa R, Ackermann M, et al. Angiogenesis in wounds treated by microdeformational wound therapy. *Ann Surg* 2011; 253: 402-9.
6. Xing D, Liu L, Marti GP, et al. Hypoxia and hypoxia-inducible factor in the burn wound. *Wound Repair Regen* 2011; 19: 205-13.
10. Zampell JC, Yan A, Avraham T, Daluvoy S, Weitman ES, Mehrara BJ. HIF-1-alpha coordinates lymphangiogenesis during wound healing and in response to inflammation. *FASEB J* 2012; 26: 1027-39.

Spleen

1. Araujo AP, Frezza TF, Allegretti SM, Giorgio S. Hypoxia, hypoxia-inducible factor-1alpha and vascular endothelial growth factor in a murine model of *Schistosoma mansoni* infection. *Exp Mol Pathol* 2010; 89: 327-33.
2. Leno-Duran E, Hatta K, Bianco J, et al. Fetal-placental hypoxia does not result from failure of spiral arterial modification in mice. *Placenta* 2010; 31: 731-7.
3. Ohta A, Diwanji R, Kini R, Subramanian M, Sitkovsky M. In vivo T cell activation in lymphoid tissues is inhibited in the oxygen-poor microenvironment. *Front Immunol* 2011; 2: 27.

Thymus

1. Hale LP, Braun RD, Gwinn WM, Greer PK, Dewhirst MW. Hypoxia in the thymus: role of oxygen tension in thymocyte survival. *Am J Physiol Heart Circ Physiol* 2002; 282: H1467-77.

2. Ohta A, Diwanji R, Kini R, Subramanian M, Sitkovsky M. In vivo T cell activation in lymphoid tissues is inhibited in the oxygen-poor microenvironment. *Front Immunol* 2011; 2: 27.

Urethra

1. Damaser MS, Whitbeck C, Chichester P, Levin RM. Effect of vaginal distension on blood flow and hypoxia of urogenital organs of the female rat. *J Appl Physiol* 2005; 98: 1884-90.

Vagina

1. Damaser MS, Whitbeck C, Chichester P, Levin RM. Effect of vaginal distension on blood flow and hypoxia of urogenital organs of the female rat. *J Appl Physiol* 2005; 98: 1884-90.