

APPLICATION OF PIMONIDAZOLE – MALIGNANT TISSUE

1. In Vitro Cells, Spheroids and Engineered Tissue Models

1. Cowan DS, Hicks KO, Wilson WR. Multicellular membranes as an in vitro model for extravascular diffusion in tumours. *Br J Cancer Suppl* 1996; 27: S28-31.
2. Sminia P, Acker H, Eikesdal HP, et al. Oxygenation and response to irradiation of organotypic multicellular spheroids of human glioma. *Anticancer Res* 2003; 23: 1461-6.
3. Prabhakaran K, Sampson DA, Hoehner JC. Neuroblastoma survival and death: an in vitro model of hypoxia and metabolic stress. *J Surg Res* 2004; 116: 288-96.
4. Chou SC, Azuma Y, Varia MA, Raleigh JA. Evidence that involucrin, a marker for differentiation, is oxygen regulated in human squamous cell carcinomas. *Br J Cancer* 2004; 90: 728-35.
5. Doege K, Heine S, Jensen I, Jelkmann W, Metzen E. Inhibition of mitochondrial respiration elevates oxygen concentration but leaves regulation of hypoxia-inducible factor (HIF) intact. *Blood* 2005; 106: 2311-7.
6. Kleiter MM, Thrall DE, Malarkey DE, et al. A comparison of oral and intravenous pimonidazole in canine tumors using intravenous CCI-103F as a control hypoxia marker. *Int J Radiat Oncol Biol Phys* 2006; 64: 592-602.
7. Meng F, Evans JW, Bhupathi D, et al. Molecular and cellular pharmacology of the hypoxia-activated prodrug TH-302. *Mol Cancer Ther* 2012; 11: 740-51.